



Artículo Especial

Artículo bilingüe – Versión inglesa

40 years of advances in Bariatric Surgery

40 años de avances en Bariatría

Aniceto Baltasar, Rafael Bou, Marcelo Bengochea, Carlos Serra, Lirios Ferri, Nieves Pérez, Francisco Arlandis, Rosa Martínez, Luis Cipagauta

Virgen de los Lirios Hospital and San Jorge Clinic, Alcoy, Spain

Abstract

Introduction: Bariatric surgical practice changes in the community setting may be under-reported. We present the developments in a Spanish bariatric surgical practice in the community setting of Alcoy from 1977 through the present.

Methods: Bariatric surgical techniques employed in a country community setting over the course of nearly four decades were reviewed retrospectively and qualitatively.

Results: Surgeons and medical professionals from Alcoy, Spain were involved in the evolution of bariatric surgery patient management and surgical technique from 1977s through 2017. During the last 40 years, 1,495 patients were treated in our clinics. Spanish bariatric surgeons contributed to advances in gastric bypass (**GBP**) in the 1970s, vertical banded gastroplasty (**VBG**) in the 1980s, biliopancreatic diversion/duodenal switch (**BPD/DS**) in the 1990s, and innovations associated with laparoscopy from the 1990s onward. Outcomes and approaches to prevention and treatment of bariatric surgical complications are reviewed from a community perspective. Contributions to the bariatric surgical nomenclature and weight-loss reporting are noted.

Conclusions: The practice of bariatric surgery in the community hospital and private clinic must be updated continuously, as in any human and surgical endeavor. Medical professionals in community bariatric practices should contribute their experiences to the field through all avenues of scientific interaction and publication.

Keywords

Obesity surgery; Bariatric surgery; Community hospital; Bariatric tendencies

Resumen

Introducción: Los cambios en la práctica de cirugía bariátrica en un hospital comarcal han sido muy importantes. Presentamos la evolución en el Hospital Comarcal de Alcoy (HCA) desde su origen en 1977 hasta el presente.


Métodos: Se revisan retrospectivamente las técnicas quirúrgicas bariátricas empleadas en un entorno comarcal a lo largo en cuatro décadas.

Resultados: Los cirujanos Alcoy, han estado involucrados en la evolución de la gestión de los pacientes de cirugía bariátrica y las técnicas quirúrgicas desde 1977 hasta la actualidad. Durante los 40 años transcurridos, 1.495 pacientes fueron tratados en nuestras clínicas comenzando con la derivación gástrica (**DG**) en la década de 1970, la gastroplastia vertical anillada (**GVA**) en la década de 1980, el cruce duodenal (**CD**) bilio-pancreático en la década de 1990, y con el acceso por vía laparoscópica desde la década de 1990.

* Autor para correspondencia.

Correo electrónico: baltasarani@gmail.com (Aniceto Baltasar).

Recibido el 15 de junio de 2017; aceptado el 24 de junio de 2017.



Los artículos publicados en esta revista se distribuyen con la licencia:
Articles published in this journal are licensed with a:
Creative Commons Attribution 4.0.
<https://creativecommons.org/licenses/by-nc-nd/4.0/>
La revista no cobra tasas por el envío de trabajos,
ni tampoco cuotas por la publicación de sus artículos.

Los resultados y los enfoques para la prevención y el tratamiento de las complicaciones de la cirugía bariátrica así como la contribución en la nomenclatura de cirugía bariátrica y la notificación de pérdida de peso son revisados desde una perspectiva comarcal.

Conclusiones: La práctica de la cirugía bariátrica en el entorno comarcal debe ser actualizada continuamente. Los cirujanos bariátricos pueden contribuir con sus experiencias en el ámbito comarcal con actualizaciones y publicaciones.

Palabras clave

Cirugía de la obesidad; La cirugía bariátrica; Hospital comarcal; Tendencias bariátricas

Abbreviations

1 ^{eu}	1 st in Europe
1 ^{sp}	1 st in Spain
1 ^w	1 st in the world
AL	Alimentary limb
ABS	Adolescent bariatric surgery
ASBS	Amer. Society for Bariatric Surgery
ASMBS	American Society for Metabolic And Bariatric Surgery
AEC	Asociación Española de Cirujanos
BMI	Body Mass Index = kg/m ²
BS	Bariatric surgery
BOLD	Bariatric Outcomes Longitudinal Data
BPD	Biliopancreatic diversion
BPD/DS	BPD/Duodenal switch
BPL	Biliopancreatic limb
CC	Common channel
CPM	Caloric-protein malnutrition
DIA	Duodenoileal anastomosis
DS	Duodenal switch
ExBMI	Expected BMI
ECEP	European Centers of Excellence Program
EGJ	Esophago-gastric junction
FBMI	Final BMI
GBP	Gastric bypass
GERD	Gastroesophageal reflux
GGF	Gastro-gastric fistula
GT	Gastric tube
IFSO	International Federation for the Surgery of Obesity and Metabolic Disorders
IBMI	Initial BMI
JIB	Jejunum Ileal bypass
JIA	Jejunum-ileum anastomosis
LDS	Laparoscopic duodenal switch
LGB	Laparoscopic gastric band
LGBP	Laparoscopic gastric bypass

MO	Morbidly obese
ECEP	Eur. Centers of Excellence Program
PE	Pulmonary embolus
REEAD	Revista Española de Enfermedades del Aparato Digestivo
RNY	Roux-en-Y
SADI	Single anastomosis duodeno-ileum
SG	Sleeve gastrectomy
SFG	Sleeve-forming gastrectomy
SECO	Sociedad Española de Cirugía Obesidad
VBG	Vertical banded gastroplasty
VLG	Vertical laparoscopic gastrectomy
WL	Weight loss
%BMIL	% BMI loss
%EBMIL	% Excess BMI loss
%EWL	% Excess weight loss

Introduction

Morbid obesity is a pandemic condition that affects humans of all countries, races and continents. The cause is multifactorial and one of the most influential factors are changes in environment and lifestyle that support consumption by humans of more calories than they utilize. When BS is employed to treat the disease of obesity, the stomach and small bowel, organs of the digestive pathway are healthy, and not the direct cause of disease. The term, BS, comes from the Greek "*baros*" meaning "weight," and "*iatrein*" means "treatment"—treatment of the broad etiology of obesity through surgery. In order to influence the weight of a patient, a bariatric surgeon can operate on the stomach to reduce caloric intake (restriction), and/or on the small bowel, to cause malabsorption of nutrients by shortening the length of the intestine (without any bowel removal). In mixed techniques, restrictive and malabsorptive components are combined. "*Metabolic Surgery*" (**MS**) represents an expansion of the concept of BS to reflect the management of metabolic conditions, such as hypercholesterolemia and/or diabetes.

When BS is used to treat obesity, the organs, the stomach and the small intestine, are healthy and are not the direct cause of the disease. The bariatric surgeon operates in the stomach to reduce caloric intake (restriction), and / or in the small intestine, to cause poor absorption of nutrients by shortening the length of the intestine useful for full absorption (bypassing a part). In mixed techniques, the restrictive and mal-absorbing components are combined.

1950s—origins of BS/MS

Henriksson ⁽¹⁾ performed the first, isolated, *JIB* in Gothenburg, Sweden in 1952 but BS field was actually initiated in 1953 at the University of Minnesota by two groups, Varco and Buchwald and Kremen and Linner ⁽²⁾.

During the latter half of the 20th century, J. Howard Payne ⁽³⁾ and other surgeons were performing *malabsorptive operations*. In 1964, Buchwald inaugurated *MS* by *diverting the distal one-third of the small bowel* to treat hypercholesterolemia ^(4,5). Edward Mason initiated the technique of the *GBP* in 1965 ⁽⁶⁾.

Methods

The aim of this *review special article* of the surgical changes made in a 40 years period in the practice of bariatric surgery at the Virgen de los Lirios Hospital and Clínica San Jorge de Alcoy, Spain

Setting

Alcoy County is located in the Alicante province of Southeastern Spain, close to the Mediterranean Sea. The population of the city is 60,000, and the county community, 140,000. Alcoy has a 300-bed National Health Service (NHS) state of the art facility, the Virgen de los Lirios district hospital, which includes an intensive care unit and operating room with three-dimensional camera laparoscopic facilities, computerized tomography and radio magnetic scans, a blood bank, standard and interventional radiology, an explant transplantation center. San Jorge Clinic is a smaller private institution where private patients were treated. All patients were treated by the same primary surgical staff.

Results

Development of bariatric surgery in Alcoy, Spain

1970s—beginnings of BS in Alcoy

On June 19, 1977, we initiated the 1st GBP in Spain, performed as recommended by Dr. Mason⁽⁶⁾ in 18 patients, an experience not published until 1980⁽⁷⁾. Much later, in 2013⁽⁸⁾, we learned that Prof. Garcia of Seville (already deceased) had performed the 1st malabsorptive JIB⁽⁹⁻¹²⁾ in 1973 as recommended by Scott⁽¹³⁾.

1980s and 1990s —vertical banded gastroplasty

Mason⁽¹⁴⁾, in 1982, published his paper on the 1^w VBG. In 1986, we published our first experiences with VBG using staplers⁽¹⁵⁾. Dr. Andreo, our radiologist, published a description of the VBG "peanut deformity"⁽¹⁶⁾.

Gastro-gastric fistula (GGF), a communication between the gastric pouch and the gastric fundus, was the main undesirable complication of the VBG. Five of our first 10 patients had this complication. While not life-threatening, the complication destroyed the intended restriction of the operation. Therefore, in 1988, our group devised the "vertical division with linear stapler and staple-line suture", separating both staple-lines with a continuous serosal suture⁽¹⁷⁾. The technique was presented in Milan, Italy in the 1st World 1988 Surgical Week that won the *Best Poster Award*. We operated on 130 patients with this technique without a single GGF. McLean et al in 1993 published later the same technique⁽¹⁸⁾.

We reported our high expectations of VBG⁽¹⁹⁾ and reviewed the same patients 5 years later, commenting that "the VBG... was a frustrating experience"^(20,21). And less than 2 years later, VBG disappeared from the bariatric armamentarium.

Baltasar et al⁽²²⁾ published in 1987 the 1st surgery for hypercholesterolemia done in Spain by diversion of the distal one-third of the small bowel and this is also the 1st Spanish publication on MS. The operation has been abandoned with the use of statins.

In 1984 RTVE (National Spanish TV) aired "JJ Soriano, más moral que el Alcoyano", in the prestigious series "Vivir cada día". It was the 1st broadcast operation of obesity surgery in Spain. It dealt with the life of a super-obese of 148 kg and IMC-52 treated by VBG at Alcoy Hospital. And it should be noted that the hospital's phones were blocked.
https://www.youtube.com/watch?v=pQtW3H9B_kg

On September 25, 1995, we assisted Dr. Favretti⁽²³⁾ of Italy in performing the 1st LGB in Spain at the Hospital La Paz, in Madrid. Carbajo⁽²⁴⁾ publicó *la 1st experience*. However, early on we abandoned use of the technique. LGB became less and less popular, first in Europe and then in the world.

1990s—duodenal switch

On March 17, 1994, we performed the 1st open DS in Spain. Our open DS series has 527 patients (5 deaths, 0.9% mortality)⁽²⁵⁻²⁷⁾. At the ASBS meeting in Seattle in June 1995, we presented a video⁽²⁸⁾ depicting the *full division vertical*

gastrectomy and suturing of the staple line. Dr. Hess⁽²⁹⁾, saw the video and began using the continuous suture, noting that, "after seeing a video of Baltasar at the annual meeting of the ASBS, we modified our procedure for the subsequent 188 cases and had only a leak". A continuous running serosal suture of the staple line prevents leaks that may occur after SG (the first part of the DS). Dr. Hess performed the 1st DS in 1988 and Dr. Marceau⁽³⁰⁾ started on June 1990 and made the 1st DS publication in 1993 and both ought to be considered the "fathers of DS".

A second contribution of our DS video was the use of the transverse supraumbilical abdominal incision (**Fig. 1a**). This approach resulted in fewer incisional hernias, 10%, compared with the midline laparotomy incision (hernia rates of 31% to 39%). The transverse incision has 2 other advantages. One is a *cosmetic*: When patients lose weight, the size of the abdominal scar is reduced to less than one-third of that achieved with the laparotomic approach (**Fig. 1b**). The second benefit is that the transverse incision facilitates post-bariatric plastic surgery, as transpositions of the navel are readily performed with a single scar just above the pubis (**Fig. 1c**). Our group later published: a) Gastric emptying⁽³¹⁾ an intermediate study on the DS^(32,33) followed by comparative studies of VBG and DS^(34,35).



Fig.1. Transverse laparotomy and scars.

In 1990, the ASBS started *Obesity Surgery (OS)*, as their official medium for reporting scholarly research and was included into the Index Medicus in 1995. This is a major accomplishment and recognition, and then became the journal of IFSO. AB entered into ASBS membership as the 1st member in 1997 and 1st IFSO in 1998 and served on the ASBS Standards Committee under the chair of George Cowan 1998-2004.

In addition to their contribution to BS by directing OS, Drs. Mervyn Deitel and George Cowan published three books and were very influential as surgeons and book publishers⁽³⁶⁻³⁹⁾. Deitel's acclaimed 1989 book inspired AB to specialize in bariatric surgery field⁽³⁶⁾.

Mid-1990s—laparoscopic BS and SECO

Wittgrove and Clark⁽⁴⁰⁻⁴²⁾ lead the transition from *laparotomic to laparoscopic BS*. We were impressed by their 1993 1st RNY LGBP experience. In early 1996, they published their second paper. In the summer of 1996, AB was asked to review their third⁽⁴²⁾ paper for *Obesity Surgery*, which summarized outcomes in 75 patients⁽⁴²⁾. AB while making two presentations at the ACS meeting in San Francisco^(43,44) visited them in San Diego and they told him that he was the 1st visiting surgeon to do so.

Our group in Alcoy performed the 1st LGBP on January 14, 1997, and reported it in the second SECO meeting in December, 1998, and then published the results of 16 cases⁽⁴⁵⁾. At the 1998-IFSO Bruges, Belgium meeting, there were only 2 surgical films on LGBP, one from Sweden by Gustavsson⁽⁴⁶⁾, and our own⁽⁴⁷⁾; these perhaps represented the 1st 2 reports on LGBP outside of the US. We concluded somewhat presciently that it was “a beautifully designed operation, with a very difficult learning curve but that *probably will become the gold standard of bariatric surgery*.”

Serra⁽⁴⁸⁾, in 1999, reported the 1st case of internal hernia after LGBP, which was cited by Higa et al in 2003⁽⁴⁹⁾. Baltasar⁽⁵⁰⁾ reported the 1st 27 LGBP cases and received the REEAD 2000 Spanish *National Surgical Award*.

Whereas, in the initial Wittgrove technique, a #33 port was used to introduce the stem of the circular stapler, AB suggested using the *stapler without a port (Fig.2)*. Later, Dr. Wittgrove⁽⁵¹⁾ acknowledged that this was a feasible and practical approach and became the accepted standard technique (Fig.3).




Fig.2. Using the circular stapler without a port.

Spanish Society for the Surgery of Obesity (SECO)

Our group *founded SECO* with 26 members in Alicante, Spain (Fig. 3) in December 1997. During the inaugural meeting, AB became the 1st President. The second meeting was held in Granada in 1998 with Drs. Capella and Álvarez Cordero as our first honorary members. In the following year, *SECO became the 13th IFSO member society*. SECO was accepted into the Spanish Surgical Society (AEC) in 2002, and AB was made *SECO Honorary President*. While AB was President of IFSO and SECO, we organized the *VIII World Congress of IFSO-2003* and the *1st Ibero-American* meetings in Salamanca. By 2004, SECO had the second highest number of publications in OS after the US, and continued to hold that position in 2005, 2006, and 2007, and held the third position in 2009⁽⁵²⁾.

SECO has been proved to employ an effective operational structure. It has had 7 presidents, with that role changing every 2 years. SECO has created sub-brands, e.g., Fun-SECO, Campus-SECO, and Multidisciplinary-SECO. Each of these groups' respective activities and information are readily accessed through its website (<http://www.seco.org/>). SECO now has 420 members and several honorary members who rank among the most distinguished bariatric surgeons in the world.

SECO Founding Members		
December 12-13, 1997		
Residencia Pérez Mateos, San Juan, Alicante, Spain		
1.	Aniceto Baltasar	Alcoy
2.	Juan Pujol	Barcelona
3.	Miguel A. Carbajo	Valladolid
4.	Santiago Tamames	Madrid
5.	Carlos Escalante	Santander
6.	Santiago Tamames	Madrid
7.	Horacio Urquijo	Madrid
8.	Cándido Martínez	Vitoria
9.	Francisco Arlandis	Alcoy
10.	Rafael Bou	Alcoy
11.	Miguel A Calvo Ros	Bilbao
12.	Antonio Alastrué	Badalona
13.	Eugenio Urquijo	Madrid
14.	Carlos Cerquella	Madrid
15.	Felipe de la Cruz	Madrid
16.	Mario García	Madrid
17.	Luis García Vallejo	Santiago
18.	Federico Leruite	Granada
19.	Juan Machuca	La Coruña
20.	José M ^a Recio	Barcelona
21.	Carlos Masdevall	Barcelona
22.	Salvador Serrano	Burgos
23.	Tomeu Feliú	Gerona
24.	Antonio Soro	Mallorca
25.	Antonio Martín	Madrid
26.	Mariano Martínez	Zaragoza

Fig.3. Founding members of Sociedad Española de Cirugía Obesidad (SECO)

Spanish bariatric surgeons are well-considered and two of them have developed techniques that are worldwide used today. Dr. Miguel Carbajo of Valladolid⁽⁵³⁾ on 2002 developed the OAGB (one anastomosis gastric bypass - BAGUA in Spanish) and Drs. Andrés Sánchez Pernaute and Antonio Torres⁽⁵⁴⁾ at Madrid Clinic Hospital the SADI (Single anastomosis duodeno-ileal).

IFSO has recently appointed Dr. Antonio Torres, a past President of SECO, as Organizer, and Dr. Baltasar, as Honorary President, of the upcoming Annual IFSO-2019 Congress in Madrid. Spain is the third country (after Austria and Brazil) to organize 2 world conferences, and the second country with 2 IFSO presidents.

Our group's members have operated as invited surgeons in Portugal (1997, 1st VBG; 2001, 1st DS; 2005, 1st SG); in Brazil (1998, Sao Paulo, 1^{er} DS; 2004, Itajai, 2nd LDS); in Buenos Aires, Argentina (2002, 1st LGBP), in Ahmedabad, India (2005, 2nd LDS), in Jerusalem, Israel (2008, 1^{er} LDS), and in Norway (2002, Forde, 1^{er} LDS; 2002, Tonsberg, 2nd LSG). We have also been guest surgeons in several national hospitals and a guest speaker in 2003 at the 1st Spring ACS Meeting, the 1st ASBS Meeting in New York, and the 1st Fall meeting of ACS-ASBS in Chicago and at the Palace of

Medical Sciences of Moscow in 2009. We have also had the privilege of hosting 71 *visiting guest surgeons* who have traveled to Alcoy, Spain to learn LDS and LSG. AB is *Honorary Member* of Brazil, Argentina and Portugal bariatric societies

Early 2000s—laparoscopic duodenal switch

Our group was present in 1999 when Gagner performed the 1st LDS in Terrassa, Spain, even before he and Ren⁽⁵⁵⁾ described it in 2000 as the most complex BS technique. We performed our 1st LDS on October 5, 2000^(56–59); Weiner et al⁽⁶⁰⁾ considered it to be the 1st LDS performed in Europe. We have done 413 LDS cases (2 deaths, 0.48%). Our LDS is performed by 3 surgeons using 6 ports (one “ #12 *single working port*” in the right upper quadrant, one #10 port for the camera, and four 5-mm working ports). All bowel divisions are accomplished with staplers and all four anastomosis are hand sewn.

In 2000, we published the 1st Spanish book in BS, “*Obesidad y Cirugía*” (*Obesity and Surgery*) with 2 editions⁽⁶¹⁾. The book was intended for the general public but was also widely read by Spanish medical professionals.

In 2005, SOARD, a journal on par with OS in the BS field, published its first volume. During these years, our publications described *gastric emptying after DS*⁽⁶²⁾, as well as *pancreato-cutaneous fistula*⁽⁶³⁾, *cardiac tamponade* by central catheter⁽⁶⁴⁾, *hepatic impairment*^(65,66), *liver failure and transplant*⁽⁶⁷⁾, *total gastrectomy*⁽⁶⁸⁾, and different types of *DS reversal*⁽⁶⁹⁾. Thirty patients (3.2%) of the 923 DS patients studied underwent *bowel lengthening* for malnutrition. We also described⁽⁷⁰⁾ the 1st occurrence of *mucosal jejunum wall hernias through the thin muscular bowel wall of patients with malnutrition* and suggested that, in malnutrition, laparotomy may be safer than laparoscopy. We performed *laparoscopic conversions from RNY to BPD* as a successful and fairly easy therapy⁽⁷¹⁾.

Mid-2000s—laparoscopic sleeve gastrectomy and other innovations

We completed our 1st LSG on December 12, 2002 and reported it 2005⁽⁷²⁾. Ahmad et al⁽⁷³⁾ recently *ranked this paper's citation index 61st within BS*. We published additional papers on LSG^(74–75) and the performance of one of the 1st *re-sleeves gastrectomies*^(76,77).

We operated LSG on 342 patients. Three of them experienced bleeding, 3 required an emergency intra-operative tracheotomy due to impossible intubation⁽⁷⁸⁾, and there were 3 EGJ leaks, and 2 deaths (0.6%) (one due to bleeding on the abdominal port and the second for EGJ leak). The entire LSG group had 93% EWL at 2 years, probably due to the systematic use of the very narrow sleeves and antrectomy, and to beginning the gastric stapling right at the pylorus⁽⁷⁹⁾.

Prevention, detection, and management of *LSG leaks* and fistulas at the EGJ have been a main concern for our group. At the IFSO-2000 meeting in Genoa, Italy we presented our use of the 1st *non-removable stent*⁽⁸⁰⁾. At that time removable-ones were not available, and so we used a non-removable Wall-stent that saved the patient's life and that lasted 6 months until esophageal hypertrophy of the mucosa caused obstruction of the esophagus requiring a total gastrectomy with intra-thoracic anastomosis⁽⁸⁸⁾. In 2007 Serra published our experience in *self-expanding stents*⁽⁸¹⁾.

Salvage surgery for chronic EGJ fistulas

In 2007 we used the 1st *RNY diversion to correct chronic EGJ fistulas*^(82–84) and presented it for discussion at the 2nd and 3rd *world LSG summits* organized by Gagner et al and Deitel et al^(85–86). *This technique has become the standard management of this serious condition (Fig. 4).*




Fig.4. RNY-diversion to treat chronic leaks by end-to-side or side-to-side.

Diabetes surgery

On February 10, 2004 we performed ^(87,88) the 1st diabetes surgery in Spain in a non-morbidly obese individual (BMI of 33 kg/m²) using the *DS without gastrectomy*, this technique was later reported in 2014 by Marceau et al ⁽⁸⁹⁾. Bou et al published our technique of *SG with duodenal-jejenum diversion (SG+DJD)* for diabetes ⁽⁹⁰⁾.

In 2006, we published a report of our 1,000 bariatric operations ⁽⁹¹⁾, which won the *Spanish National Surgical Award*.

Adolescent BS (ABS)

A 10-year-old wheelchair-bound boy with a BMI of 42 kg/m², Blount's disease, and knee fractures underwent LSG in 2008 ⁽⁹²⁾. The patient was the 1st ABS done in Spain, and one of the world youngest BS patients at the time. He is doing very well nine years later ⁽⁹³⁾, and has no impairment in his growth with a BMI of 26 Kg/m².

European centers of excellence program

In 2009, Drs. Scopinaro, Melissas, Fried, and Baltasar created the IFSO European Chapter of the Centers of Excellence (COE) Program ⁽⁹⁴⁾. Multiple European centers and surgeons utilize this program currently.

2010s—Publications and technical improvements

BMI-Latina (Bariatric and Metabolic Ibero-Americana) was founded by us in 2011 as online journal published in Spanish, English, and Portuguese. The magazine was adopted by SECO in 2015 as the Spanish society's magazine and we believe it will play a valuable role in BS in the Spanish and Portuguese speaking societies but since from 2015 only the abstracts of the English edition are published currently ⁽⁹⁵⁻⁹⁸⁾.

Recently, Serra et al ⁽⁹⁹⁾ recommended 2 new technical improvements in SG, including our 1st publication on "sliding self-locking stitch" and the Aberdeen knot. Baltasar ⁽¹⁰⁰⁾ described its use and he recommends the *antrectomy starting at the pylorus plus an omental patch* to cover the staple line and prevent rotation of the sleeve. We also called attention to the need to *prevent stapling of the bougie* ⁽¹⁰¹⁾ as well as a simple Whaps up bariatric telemedicine solution to detect early complications after hospital discharge ^(102,103).

By 2013, Spain was the country with the fifth most publications in SOARD. Now, in 2017, our group has 119 BS publications; we have reviewed >74 papers for *Obesity Surgery*, and >14 for SOARD, and other national and international journals. At the IFSO 2009 meeting in Paris, AB received the *IFSO Lifetime Membership Award*. On June 2011, at the ASMBS meeting in Orlando, Florida, AB ⁽¹⁰⁴⁾ was second as a *finalist for the prestigious ASMBS Outstanding Achievement Award* (this is the "Oscar" of BS).

Discussion

In addition to participating in the aforementioned technical changes in the BS field, our Alcoy practice has contributed commentary on key concepts in the BS nomenclature and on reporting weight loss ⁽¹⁰⁵⁾.

Nomenclature

BS is a relatively new specialty. We have long been interested in the development of BS nomenclature ⁽¹⁰⁶⁻¹⁰⁷⁾; our group, under the direction of the Royal Spanish Academy of Medicine, agreed that the correct name in Spanish for the SG technique is "*vertical laparoscopic gastrectomy*" (**VLG**) ⁽¹⁰⁸⁾, and also, that the end result of the VLG is the formation of a "*gastric tube*," or "*sleeve*"—and so it was approved in the General Assembly of the 2010 SECO meeting in Valencia.

We also published that the English the term, "*SG is an inappropriate, a misnomer term*" ⁽¹⁰⁹⁾ as the meaning of "gastrectomy" is "amputation and elimination" of any part of the stomach. As right colectomy means the elimination of the right colon, not the left, and a left nephrectomy is the removal of the left and not the right kidney, the use of the term SG should more precisely mean "withdrawal of the sleeve," although, in fact, the sleeve is not removed, it is the sleeve portion of the stomach that remains. Therefore, we have suggested that the SG name is a *misnomer* and the correct term might be "**sleeve-forming gastrectomy**" (**SFG**) ⁽¹¹⁰⁾, meaning more precisely a gastrectomy that forms a sleeve. We recognize that the term SG is so ingrained in common usage that it will likely prevail. Surgeons, besides performing excellent procedures, should also aim to speak and write correctly.

Weight-loss reporting

Weight-loss reporting is also controversial in BS. We have suggested that BMI measurement is much better than using WL measurement since it takes into account a patient's height ^(111,102). We should not expect all patients with such different IBMIs to reach an FBMI close to the ideal BMI of 25 kg/m². This is even more the case for high IBMI patients. Dr. Borrás, our group's mathematician, devised a mathematical system to measure the "*expected BMI*" = *ExBMI* = goal BMI, by using 7,410 real patients from multiple centers ⁽¹¹³⁾. The ExBMI mathematical formula for each patient is $FBMI = IBMI \times 0.4 + 12$. By using the BOLD system with a large patient sample, it will be possible to apply a better final mathematical formula using the concept of ExBMI. Using the ideal %ExBMI of 100%, any FBMI above that may be considered a success, and that below, a failure. Grading of the results depends on the extent of the deviation. A *preliminary report* of WL in our DS cases shows that at 5, 10, and 15 years, BMIL of 85%, and an ExBMI of 96% suggest that almost all patients had an FBMI close to the goal results. Molina has recently confirmed these results ⁽¹¹⁴⁾.

BMI should be the *standard measurement* of obesity. DS is the most effective BS operation because it combines reduction on the intake and malabsorption as shown in the Fig. 5. A patient if with a BMI-100 the *largest ever operated in Spain* that dropped to BMI-34.

In total, 1494 patients with obesity were treated. The duodenal switch was used in 987 (515 ODS 472 LDS) with 0.7 mortality, one of the largest series in the world (of which 376 were from USA and Canada, 73 from Norway and 25 from England were operated on at the San Jorge Clinic)

Our videos about obesity can be seen at

www.youtube.com/channel/UCepwVhQEohunxKOhNqC71zg/videos?view=0&sort=dd&shelf_id=0



BMI-100, 231 Kg, 149 cm

6 months - 132 Kg

2 years- 79 Kg, FBMI-34

102 Kg lost

153 Kg Lost

Fig 5. BMI changes in a super-super obesity DS from 100 to 34

Conclusion

For the past 4 decades, from our community practice setting, we have followed the changes in the practice of BS throughout the world and participated in the evolution of the field. It has been extremely important and satisfying to continuously share and discuss evidence with our peers through the scientific literature, and in person, in local, national, and international interactions and meetings. We believe that small institutions can provide high-quality bariatric surgical care to patients and make significant scientific contributions to the field.

Acknowledgments

We would like to thank the editors and all the cited colleagues in this paper for their enormous help, and for the pleasure we have received in meeting and coming to know them all, and also to learn from their excellent teachings.

Ethical Approval

All procedures involving human participants were performed in the studies cited herein in accordance with the ethical standards of the institutional and national research committees and with the Helsinki Declaration and amendments.

Informed Consent

Informed consent was obtained from all individual participants included in the studies cited.

Conflict of Interest

None of the authors has any conflicts of interest to disclose. J. N. Buchwald, Medwrite Medical Communications, WI, USA, received a fee for substantive manuscript revision.

References

1. Henrikson V. Is small bowel resection justified as treatment for obesity? *Nordisk Medicin* 1952; 47: 744
2. Linner JH. Overview of surgical techniques for the treatment of morbid obesity. *Gastroenterol Clin North Am* 1987; 16:253-72.

3. Payne JH, De Wing L, Commons RR: Metabolic Observations in Patients with Jejunocolic Shunts. *Am J Surg* 1963; 106:273-89
4. Buchwald H. Lowering of the cholesterol absorption and blood levels by ileal exclusion. *Circulation*.1964; 29:711-50.
5. Shikora SS. Biography: Henry Buchwald, M.D., Ph.D. *Obes Surg*. 2016;26:3-4.
6. Mason EE, Ito C. Gastric bypass in Obesity. *Surg Clin North*. 1967;47:1345-52.
7. Baltasar A, Del Río J, Bengochea M. Bypass gástrico en la obesidad mórbida. *Rev Clín Española*.1980; 157:395-9.
8. Baltasar A. Dominguez-Adame. Beginnings of bariatric and metabolic surgery in Spain. *Cir Esp*.2013;91:413-6.
9. García Díaz S. Indicaciones del tratamiento quirúrgico en la obesidad. *Rev Quir Esp*. 1975;2:165-72.
10. García S, Pera C, Loscertales J, Arévalo E. Estado actual de la cirugía de la obesidad. *REEAD*. 1977; 51:1-25.
11. García S, Pera C, Loscertales J, Arévalo JE. Cirugía de la obesidad. *Advances Cirugía*. 1979; 2:209-24.
12. García S, García G. Medical and surgical indications for treatment of morbid obesity. *World J Surg*. 1981; 5:795-9.
13. Scott H, Dean R, Skull J et al. New considerations in use of yeyuno-ileal by-pass in morbid obesity. *Ann. Surg*. 1973; 177:723
14. Mason EE. Vertical banded gastroplasty in obesity. *Arch Surg*. 1982; 157:701-6.
15. Baltasar A. Suturas mecánicas en la cirugía de la obesidad. *Rev Soc Andaluza Patología Digestiva*. 1987; 10:555-6.
16. Andreo L, Lasierra R, Baltasar A, et al. Evaluación radiológica de la gastroplastia vertical anillada. *Radiología*. 1988;30:303-8.
17. Baltasar A. Modified vertical gastroplasty: Technique with vertical division and serosal patch. *Acta Chir Scand*. 1989;155:107-12.
18. McLean LD, Rhode BM, Sampalis J, et al. Results of the surgical treatment of obesity. *Am J Surg*. 1993; 165:155-62.
19. Baltasar A, Tomás J, Marcote E, et al. Cirugía bariátrica: Experiencia con 100 casos operados. *Cir Esp*. 1991;50:271-7.
20. Baltasar A, Bou R, del Río J, et al. Cirugía bariátrica: Resultados a largo plazo de la gastroplastia vertical anillada. ¿Una esperanza frustrada? *Cir Esp*. 1997;62:175-9.
21. Baltasar A, Bou R, Arlandis F, Martínez R, et al. Vertical banded gastroplasty at more than five years. *Obes Surg*. 1998;8:29-34.
22. Baltasar A, Marcote E, Bou R, et al. Exclusión ileal parcial en la hipercolesterinemia. *Cir Esp*. 1991;49:475-8.
23. Favretti F, Cadière G.B, Segal G.: Laparoscopic Banding: Selection and Technique in 830 Patients Obesity Surgery,12, 385-39
24. Carbajo MA, Martin JC, Blanco JJ.: Banda gástrica ajustable en el tratamiento de loa obesidad mórbida. *Cir Esp* 1988, 63:185-8
25. Baltasar A, Bou R, Marcote E, et al. Hybrid bariatric surgery: Biliopancreatic diversion and duodenal switch. *Obes Surg*. 1995;5:419-23.
26. Baltasar A. El cruce duodenal: Una terapia efectiva en la obesidad mórbida. Estudio intermedio. *Cir Esp*. 2001;69:445-50.
27. Baltasar A. Preliminary results of the duodenal switch. *Obes Surg*. 1997;7:500-4.
28. Baltasar A, Bou R, Bengochea M, et al. Cirugía híbrida bariátrica: Cruce duodenal en la derivación Biliopancreática por obesidad. *VRC*. 1996;12:16-41. www.youtube.com/watch?v=h0nTzeUDI5o
29. Hess DS, Hess DW. Biliopancreatic diversion with a duodenal switch. *Obes Surg*. 1988;8:267-82.
30. Marceau P, Biron S, Bourque RA et al: Biliopancreatic Diversion with a New Type of Gastrectomy. *Obesity Surgery* 1993,3:29-35.
31. Martínez RM, Baltasar A. Estudio del vaciado gástrico en pacientes con obesidad mórbida intervenidos por cruce duodenal. *Rev Esp Enf Digest*. 1997;89:413-4.
32. Baltasar A, Bou R, Bengochea M, Arlandis F. et al. Duodenal switch: An effective treatment for morbid obesity. Intermediate study. *Obes Surg*. 2001;11:54-9.
33. Baltasar A. El cruce duodenal: Una terapia efectiva en la obesidad mórbida. Estudio intermedio. *Cir Esp*. 2001;69:445-50.
34. Pérez N, Baltasar A, Serra C, et al. Comparative analysis of vertical banded gastroplasty and duodenal switch at five years follow-up. *Obes Surg*. 2005;15(7):1061-5.
35. Pérez N, Serra C, Baltasar A, et al. Resultado comparativo entre la GVA y el CD en la obesidad mórbida. *Cir Esp*. 2006;79:283-8.
36. Deitel M. Surgery for the morbidly obese patient. London: Lea & Febiger, 1989.

37. Deitel M, Shikora S. Mervyn Deitel—Biography. *Obes Surg.* 2016;26:707–8.
38. Deitel M, Gagner M, Dixon JB, Himpens J, Madan A (eds). *Handbook of Obesity Surgery.* Toronto: FD-Communications Inc., 2010.
39. Deitel M, Cowan G. Update: Surgery for the Morbidly Obese Patient 2000. Toronto: FD-Communications, Inc., 2000.
40. Wittgrove A, Clark W, Tremblay L. Laparoscopic gastric bypass. Roux-en-Y. Preliminary report. *Obes Surg.* 2004;4:353–7.
41. Wittgrove A, Clark W. Laparoscopic gastric bypass. Roux-en Y. Experience with 27 cases with 3-18 months follow up. *Obes Surg.* 2006;6:54–7.
42. Wittgrove A, Clark W, Schubert K. Laparoscopic gastric bypass. Roux-en Y. Technique and results in 75 patients with 3-30 months follow up. *Obes Surg.* 2006;6:500–4.
43. Martínez R, Arlandis F, Baltasar B, et al. Vascular (aorta) reconstruction for retroperitoneal sarcomas I, VRC. 1999;16.4:15–7.
44. Baltasar A. Iterative oncologic-vascular surgery for recurrent retroperitoneal sarcoma. Vascular trans-pubis cross-over (arterial and venous) re-construction for retroperitoneal sarcomas II. VRC. 1999;16.4:31–4.
45. Baltasar A, Bou R, Miró J, Serra C, et al. Bypass gástrico por laparoscopia. *Cir Esp.* 2000;67:556–9.
46. Gusfavnson S, Westling A. Laparoscopic Roux-en-Y gastric bypass. Uppsala, Sweden. *Obes Surg.* 1998;8:392.
47. Baltasar A, Bou R, Bengochea M, Miró J, Arlandis F. Lap Roux-Y gastric bypass. Alcoy. Spain. *Obes Surg.* 1998;8:393.
48. Serra C, Baltasar A. Internal hernias and gastric perforation after laparoscopic gastric bypass. *Obes Surg.* 1999;9:546–9.
49. Higa K, Ho T, Boone K. Internal hernias after laparoscopic Roux-en-Y gastric bypass: Incidence, treatment and prevention. *Obes Surg.* 2003;13:350–4.
50. Baltasar A, Bou R, Miró J, et al. Laparoscopic gastric bypass. Bypass gástrico laparoscópico. *Rev Esp Enferm Ap Dig.* 2000;10:661–4.
51. Wittgrove A. Chapter 21: Laparoscopic gastric bypass. Circular stapler technique. In: *Obesity Surgery: Principles and Practice* (eds. Pitombo C, Jones KB, Higa KD, Pareja JC, eds). Albany, New York: McGraw Hill Medical;2007, 221-224.
52. Dabi Y, Darrigues L, Katsahian S et al.: Publication Trends in Bariatric Surgery: a Bibliometric Study. *Obes Surg* 2016; 26:2691–2699. DOI 10.1007/s11695-016-2160-x
53. Carbajo M, García-Caballero M, Toledano M et al : One-Anastomosis Gastric Bypass by Laparoscopy: Results of the First 209 Patients. *Obesity Surgery* 2005, 15, 398-404
54. Pernaute A, Perez E, Torres et al: Modern Surgery: Technical Innovation “Right-Angled” Stapled Latero-lateral Duodeno-jejunal Anastomosis in the Duodenal Switch. *Obesity Surgery*2005, 15, 700-702
55. Ren CJ, Patterson E, Gagner M. Early results of laparoscopic biliopancreatic diversion with duodenal switch: a case series of 40 consecutive patients. *Obes Surg.* 2000;10:514–23.
56. Baltasar A. Bou R, Miro J, et al. Laparoscopic duodenal switch. Technique and initial experience. *Obes Surg.* 2001;12:245–8.
57. Baltasar A. Hand-sewn laparoscopic duodenal switch. *Surg Obes Relat Dis.* 2007;3:94–6.
58. Baltasar A. Chapter 29: Laparoscopic biliopancreatic diversion. The duodenal switch. In: *Obesity Surgery: Principles and Practice* (eds. Pitombo C, Jones KB, Higa KD, Pareja JC, eds). Albany, New York: McGraw Hill Medical; 2007, pp.241–3.
59. Baltasar A. El Cruce Duodenal. *Guía Clínica de obesidad de Cirugía Española*, Chapter 29. Madrid, Spain: S.A. Arán Ediciones, 2007.
60. Weiner RA, Blanco-Engert R, Weiner S, et al. Laparoscopic biliopancreatic diversion with duodenal switch: Three different duodeno-ileal anastomotic techniques and initial experience. *Obes Surg.* 2004;14:334–40.
61. Baltasar A. *Obesidad y Cirugía. Cómo dejar de ser obeso.* Madrid, Spain: S. A. Aran Ediciones, 2000.
62. Martínez R, Baltasar A, Vidal V, et al. Estudio de vaciado gástrico en pacientes con obesidad mórbida intervenidos por cruce duodenal. *REED.* 1997;89:413–4.
63. Bueno J, Pérez N, Serra C, Bou R, Miró J, Baltasar A. Fístula pancreato-cutánea secundaria a pancreatitis postoperatoria tras cruce duodenal laparoscópico. *Cir Esp.* 2004;76(3):184–6.
64. Serra C, Baltasar A, Losa JL, et al. Taponamiento cardiaco provocado por perforación de catéter venosa central. *Cir Esp.* 1999;65:470–1.
65. Baltasar A, Serra C, Pérez N, et al. Clinical Hepatic Impairment after the Duodenal Switch. *Obes Surg.* 2014;14:77–8.
66. Baltasar A. Liver cirrhosis and bariatric operations. *Surg Obes Relat Dis.* 2006;2:580–1.

67. Baltasar A. Liver failure and transplantation after duodenal switch. *Surg Obes Relat Dis.* 2014;10:e93–6.
68. Serra C, Baltasar A, Pérez N, et al. Total gastrectomy for complications of the duodenal switch. *Obes Surg.* 2006;16:1082–6.
69. Baltasar A, Bou R, Bengochea M, Serra C. Protein caloric malnutrition. Three types of bowel lengthening. *BMI-Latina.* 2011;1.2.5:96–7.
70. Baltasar A, Bou R, Bengochea M. Fatal perforations in laparoscopic bowel lengthening operations for malnutrition. *Surg Obes Relat Dis.* 2010;20:140–7.
71. Baltasar A. Letter to the editor: Roux-en-Y to duodenal switch. *Surg Obes Relat Dis.* 2008;4:208–11.
72. Baltasar A, Serra C, Pérez N, et al. Laparoscopic sleeve gastrectomy. A multi-purpose operation. *Obes Surg.* 2005;15:1124–8.
73. Ahmad SS, Ahmad SS, Kohl S. The Hundred Most Cited Articles in Bariatric Surgery. *Obes Surg.* 2015;25:900–9.
74. Serra C, Pérez N, Bou R, Bengochea M, Martínez R, Baltasar B. Gastrectomía tubular laparoscópica. Una operación bariátrica con diferentes indicaciones. *Cir Esp.* 2006;79:289–92.
75. Sánchez-Santos R, Masdevall C, Baltasar A, et al. Short- and mid-term outcomes of sleeve gastrectomy for morbid obesity: The experience of the Spanish National Registry. *Obes Surg.* 2009;19:2019–23.
76. Baltasar A, Serra C, Pérez N. Re-sleeve gastrectomy. *Obes Surg.* 2006;16:1535–8.
77. Serra C, Baltasar A. Re-gastrectomía tubular. *Cir Esp.* 2007;82.1:240.
78. Baltasar A, Bou R, Bengochea M, Serra C, Pérez N. Intubación difícil y traqueotomía en el obeso mórbido. Difficult intubation and tracheotomy in the morbidly obese. *BMI-Latina.* 2013;3:4–7.
79. Baltasar A, Bou R, Bengochea M, Serra C, Pérez N. Laparoscopic gastric sleeve, subtotal antrectomy and omentoplasty. *Obes Surg.* 2015;25:195–6.
80. Baltasar A, Pérez N, Bou R, Miro J, Bengochea M, Arlandis F, Escrivá C, Serra C, Martínez R. Wall-stent prosthesis for severe leak and obstruction of the duodenal switch Gastric tube. *Obes Surg.* 2000;10:309.
81. Serra C, Baltasar A, Andreo L, et al. Treatment of gastric leaks with coated self-expanding stents after sleeve gastrectomy. *Obes Surg.* 2007;17:866–2.
82. Baltasar A, Bou R, Bengochea M, et al. Use of a Roux limb to correct esophagogastric junction fistulas after sleeve gastrectomy. *Obes Surg.* 2007;17:1409–10.
83. Baltasar A, Serra C, Bengochea M, Bou R. The use of a Roux-en-Y loop for sleeve gastrectomy fistulas. *Bariátrica & Metabólica.* 2008;3:20–3.
84. Baltasar A, Serra C, Bengochea R, et al. Use of Roux limb as remedial surgery for sleeve gastrectomy fistulas. *Surg Obes Relat Dis.* 2009;4:759–63.
85. Gagner M, Deitel M, Traci L, et al. The second international consensus summit for sleeve gastrectomy, March 19–21, 2009. *Surg Obes Relat Dis.* 2009;5:476–85.
86. Deitel M, Gagner M, Erickson A. et al. Third international summit: current status of sleeve gastrectomy. *Surg Obes Relat Dis.* 2011;7:749–59.
87. Baltasar A. Historical note: First diabetes metabolic operation in Spain. *Int Obes Dia.* 2015;25:180–2.
88. Baltasar A. Primera operación de diabetes en España. *BMI-Latina.* 2016;6:996–9.
89. Marceau R, Biron S, Marceau S et al. Biliopancreatic diversion-duodenal switch: Independent contributions of sleeve resection and duodenal exclusion. *Obes Surg.* 2014;24:1843–9.
90. Bou R, Baltasar A, Pérez N, et al. Cirugía de la diabetes: Gastrectomía vertical y desvío duodenal. *BMI-Latina.* 2012;2:400–3.
91. Baltasar A, Bou R, Bengochea M, Serra C, Pérez N. Mil operaciones bariátricas. One thousand bariatric operations. *Cir Esp.* 2006;79:349–55.
92. Baltasar A, Serra C, Bou R, et al. Sleeve gastrectomy in a 10-year-old child. *Obes Surg.* 2008;18:33–6.
93. Baltasar A. Nine years follow up of the 1st Adolescent Bariatric Surgery in Spain. *J J Obesity.* 2017. 3(1): 026
94. Scopinaro N, Melissas J, Fried M, Baltasar A. European Chapter of Centers of Excellence Program. Web: <http://www.eacbs.com/site/index.php/ifso-endorsed-coe-programs/ifso-ec-coe-program>
95. Baltasar A. Editorial. *BMI-Latina.* 2011;1:5–9.

96. Baltasar A, Serra C, Bou R, et al. Complicaciones de la cirugía bariátrica. of Bariatric Surgery complications. BMI-Latina. 2011;1:238–42.
97. Baltasar A, Bou R, Bengochea M. Cruce duodenal abierto. Open duodenal switch. Video. BMI-Latina. 2011;1:357–60.
98. Baltasar A, Bou R, Bengochea M, et al. Cruce duodenal manual laparoscópico. Laparoscopic hand-sewn duodenal switch. Video. BMI-Latina. 2012;2:387–9.
99. Serra C, Pérez N, Bou R, Baltasar A. Sliding self-locking first stitch and Aberdeen Knot in suture reinforcement with omentoplasty of the laparoscopic gastric sleeve staple line. *Obes Surg.* 2014;24:1739–40. Video: https://youtu.be/MXf0aiWp_kE
100. Baltasar A, Bou R, Serra C, Bengochea M, Pérez N. Use of self-locking knots in running intestinal bariatric sutures. *Global Surg* 2015;2:100-1.
101. Baltasar A, Bou R, Serra C, et al. Video. Stapling of the bougie in the sleeve gastrectomy. *Surg Obes Relat Dis.* 2016; 12:1136-1138. DOI <http://dx.doi.org/10.1016/j.sorad.2016.01.02>.
102. Baltasar A. Simple telemedicina bariátrica. Simple bariatric telemedicine. BMI-Latina. 2016;5:900–1.
103. Baltasar A.: WhatsApp© Assistance in Bariatric Surgery. *Journal of obesity and eating disorders* 2017.3:28-9.DOI: 10.21767/2471-8513.100017
104. Shikora S, Torres A. Biography: Aniceto Baltasar. *Obes Surg* 2016;26:477–8.
105. Baltasar A. Nomenclatura utilizada en cirugía bariátrica. *Cir Esp.* 2008;83:220–1.
106. Baltasar A. Gastroplastia o gastrectomía, gastroplasty versus gastrectomy. *Cir Esp.*2009;85:325–6.
107. Baltasar A. Editorial: Nomenclatura utilizada en cirugía bariátrica. Nomenclature of bariatric terms. BMI-Latina. 2011;1:278.
108. Baltasar A. La Real Academia Nacional de Medicina de España dice. La gastrectomía vertical es el término correcto. BMI-Latina. 2012;2:381–3.
109. Baltasar A. Laparoscopic sleeve gastrectomy is a misnomer. *Surg Obes Relat Dis.*2012;8:127–31.
110. Baltasar A. Sleeve-forming gastrectomy is the right terminology. *Obes Surg.*2015;25:935–7.
111. Baltasar A, Deitel M, Greenstein R. Weight loss reporting. *Obes Surg.* 2008;18:761–2.
112. Baltasar A et al. IMC esperable tras cirugía bariátrica. Expected BMI after bariatric operations. *Cir Esp.*2009;86:308–12.
113. Baltasar A, Pérez N, Serra C, Bou R, Bengochea M, Borrás F. Weight loss reporting: Predicted BMI after bariatric surgery. *Obes Surg.* 2011;212:367–72.
114. Molina A, Fatima F, Vives M. et al.: Usefulness of Baltasar's expected body mass index as an indicator of bariatric weight loss surgery. *Obes Surg.* 2016. DOI 10.1007/s11695-016-2163-7